

## Algorithms of Scientific Computing II

### Exercise 2 - Modelling

#### 1) Pair Potentials and Forces

There are lots of different potentials describing the interaction between two entities. Examples are the harmonic potential for two bodies which are connected by a spring, the gravitational potential for any pair of objects in our universe and others. For this exercise, you will need the following potentials:

- Hard sphere potential:  $U_{HS}(r) = \begin{cases} \infty & \forall r \leq d \\ 0 & \forall r > d \end{cases}$
- Soft sphere potential:  $U_{SS}(r) = \epsilon \left(\frac{\sigma}{r}\right)^n$
- Van der Waals potential:  $U_W(r) = -4\epsilon \left(\frac{\sigma}{r}\right)^6$
- Lennard-Jones potential:  $U_{LJ}(r) = \alpha\epsilon \left(\left(\frac{\sigma}{r}\right)^n - \left(\frac{\sigma}{r}\right)^m\right)$

- a) From the formula for the pair potential, the force which acts upon the two bodies can be derived. Calculate the force for the given potentials.

**ANSWER:**

Name	potential	force	repulsive (+) / attractive(-)
Hard Sphere	$\begin{cases} \infty & \forall r \leq d \\ 0 & \forall r > d \end{cases}$	$\begin{cases} 0 & r \neq d \\ \infty & r = d \end{cases}$	+
Soft Sphere	$\epsilon \cdot \left(\frac{\sigma}{r}\right)^n$	$\frac{n \cdot \epsilon}{r} \cdot \left(\frac{\sigma}{r}\right)^n$	+
Van der Waals	$-4\epsilon \cdot \left(\frac{\sigma}{r}\right)^6$	$\frac{-24\epsilon}{r} \cdot \left(\frac{\sigma}{r}\right)^6$	-
Lennard-Jones-12-6	$4\epsilon \cdot \left(\left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6\right)$	$\frac{24\epsilon}{r} \cdot \left(2 \cdot \left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6\right)$	+ -

- b) Draw an approximate graph of all potentials and forces.

c) Examine the calculated force functions and try to find qualitative differences between them. Consider especially the following properties:

- attraction or repulsion
- influence of the distance
- usability on a computer

**ANSWER:**

- Hard-Sphere: not integratable, therefore not usable on a computer
- Modell attractive / repulsive forces or both
- One thing in common: force and potential decrease very quickly with increasing distance, therefore called short-range potentials (as opposed to long-range potentials like gravitation)

## 2) Multi-Centered Molecules

For single-centered molecules, the force on molecule  $i$  equals the sum of all forces between molecule  $i$  and all other molecules:  $\vec{F}_i = \sum_{j \neq i} \vec{F}_{ij}$

Using the force, the acceleration of molecule  $i$  is given by the following formula:

$$\ddot{\vec{x}}_i = \frac{\vec{F}_i}{m_i} = \frac{\sum_{j \neq i} \vec{F}_{ij}}{m_i}$$

Now consider multi-centered molecules. There are some more values to be considered to be able to represent rotations:


- values already considered for single-centered molecules: force  $\vec{F}$ , mass  $m$ , acceleration  $\ddot{\vec{x}}$ .
- values only to be considered for multi-centered molecules: torque  $T$ , moment of inertia  $I$ , angular acceleration  $\ddot{\omega}$ .

a) Find the formula for the angular acceleration that is analogue to the formula for the acceleration  $\ddot{\vec{x}}$ .

**ANSWER:**

A multi-centered molecule is composed of multiple centers (sites). The force between two molecules  $i$  and  $j$  on the center of molecule is calculated as the sum of the pairwise forces between the single sites:

$$F_{ij} = \sum_{sites_i} \sum_{sites_j} F_{sites}$$



The forces on the sites cause a torque on the molecule. The torque resulting from one site can be calculated as

$$\vec{\tau} = (\vec{r}_{site} - \vec{r}_{center}) \times \vec{F}_{site}.$$

where  $\vec{r}_{site}$  denotes the position of the site and  $\vec{r}_{center}$  is the position of the center of the molecule.

Thus the torque on the whole molecule  $i$  can be calculated as

$$\vec{\tau}_i = \sum_{sites} (\vec{r}_{site} - \vec{r}_{center}) \times \vec{F}_{site}$$

The relation between torque  $\tau$ , angular acceleration  $\frac{\delta\omega}{\delta t}$  and moment of inertia  $\theta$  is given by

$$\vec{\tau} = \vec{\theta} \cdot \frac{\delta\omega}{\delta t}$$

Consequently, the angular acceleration for molecule  $i$  can be computed as

$$\frac{\delta\omega_i}{\delta t} = \frac{\vec{\tau}_i}{\vec{\theta}_i} = \frac{\sum_{sites} ((\vec{r}_{site} - \vec{r}_{center}) \times \vec{F}_{site})}{\vec{\theta}_i}$$

### 3) Dimensionless formulation

a) What is the rationale of dimensionless formulations?

**ANSWER:**

- Reduce the drastic difference in the order of magnitudes of the values
- Make a statement for a class of materials (as under certain conditions, it's sufficient to perform a calculation for a given parameter set, e.g.  $\epsilon$  and  $\sigma$  for the Lennard-Jones potential. The result can then be devolved to other parameter combinations).

**b)** In order to simulate the inert gas Argon we have to create an initial configuration. The length of a timestep is  $2.17 \text{ fs}$ . The simulation domain is a cube of sidelength  $1 \mu\text{m}$ . Initially, the gas should be subject to normal conditions, i.e. the pressure is 1 bar at a temperature of  $273,5 \text{ K}$ . Further we assume, that all atoms have the same velocity.

Calculate the following values (give the values which are marked with (\*) dimensionless!):

- N: Number of atoms within the domain
- L\*: side length of the domain
- dt\*: timestep length
- v\*: velocity of an atom
- T\*: temperature

### ANSWER:

For the following calculation we use  $\sigma$ ,  $\epsilon$  and the mass of argon:

$$\begin{aligned}\sigma &= 3.41 \cdot 10^{-10} \text{ m} \\ \epsilon &= 119.8 \cdot 1.38066 \cdot 10^{-23} \text{ J} \\ &= 1.654 \cdot 10^{-21} \text{ J}\end{aligned}$$

The molar mass of argon is  $39.948 \frac{\text{g}}{\text{mol}} = 3.9948 \cdot 10^{-2} \frac{\text{kg}}{\text{mol}}$  and a mol of gas under normal conditions contains  $6.0221415 \cdot 10^{23}$  molecules, (Avogadro constant), therefore follows for the mass of an atom:

$$\begin{aligned}m &= \frac{3.9948 \cdot 10^{-2}}{6.0221415 \cdot 10^{23}} \text{ kg} \\ &= 6.6335 \cdot 10^{-26} \text{ kg}\end{aligned}$$

The Loschmidt number corresponds to the number of molecules of an ideal gas under normal conditions contained in  $1 \text{ cm}^3$ . The volume of the simulation domain is  $(1 \mu\text{m})^3 = 10^{-12} \text{ cm}^3$ .

Thus the number  $N$  of atoms is:

$$N = 2.687 \cdot 10^{19} \frac{1}{\text{cm}^3} \cdot 10^{-12} \text{cm}^3 = 2.687 \cdot 10^7$$

With the formulas for dimensionless position and time we obtain

$$\begin{aligned} L^* &= \frac{1}{\sigma} \cdot L = \frac{1}{3.41 \cdot 10^{-10} \text{m}} \cdot 10^{-6} \text{m} \\ &= 2932.6 \\ dt^* &= \frac{1}{\sigma} \cdot \sqrt{\frac{\epsilon}{m}} \cdot dt = \frac{1}{3.41 \cdot 10^{-10} \text{m}} \cdot \sqrt{\frac{1.654 \cdot 10^{-21} \text{J}}{6.6335 \cdot 10^{-26} \text{kg}}} \cdot 2.17 \cdot 10^{-15} \text{s} \\ &= 0.001 \end{aligned}$$

Bevor calculating the dimensionless formulation of the velocity, we have to calculate the “real” velocity first. The temperature of matter is defined by the kinetic energy of its molecules. The kinetic energy can be calculated from the mass and velocity of the atoms:

$$\begin{aligned} T &= \frac{2}{3 \cdot N \cdot k_B} \cdot E_{kin} \\ E_{kin} &= \frac{1}{2} \cdot \sum_i m_i \cdot v_i^2 \\ &= \frac{1}{2} \cdot N \cdot m \cdot v^2 \end{aligned}$$

Those formulas can be reformulated:

$$\begin{aligned} E_{kin} &= \frac{3}{2} \cdot N \cdot k_B \cdot T = \frac{3}{2} \cdot 2.687 \cdot 10^7 \cdot 1.38066 \cdot 10^{-23} \frac{\text{J}}{\text{K}} \cdot 273.15 \text{K} \\ &= 1.52 \cdot 10^{-13} \text{J} \\ v &= \sqrt{\frac{2 \cdot E_{kin}}{N \cdot m}} \\ &= \sqrt{\frac{2 \cdot 1.52 \cdot 10^{-13} \text{J}}{2.687 \cdot 10^7 \cdot 6.6335 \cdot 10^{-26} \text{kg}}} \\ &= 412.98 \frac{\text{m}}{\text{s}} \end{aligned}$$

From that we can calculate the dimensionless form of the velocity:

$$\begin{aligned}v^* &= \frac{dt}{\sigma} \cdot v = \frac{2.17 \cdot 10^{-15} s}{3.41 \cdot 10^{-10} m} \cdot 412.98 \frac{m}{s} \\ &= 0.0026281\end{aligned}$$

Last we have to determine the dimensionless value for the temperature:

$$\begin{aligned}T^* &= T \cdot \frac{k_B}{\epsilon} = 273.15 K \cdot \frac{1.38066 \cdot 10^{-23} \frac{J}{K}}{1.654 \cdot 10^{-21} J} \\ &= 2.2801\end{aligned}$$