


# Algorithms of Scientific Computing (Algorithmen des Wissenschaftlichen Rechnens) Adaptive Sparse Grids, Orthogonality

## 1 Adaptive Sparse Grids

Here, the exercise is to adaptively refine a 2-dimensional sparse grid without boundary. We follow the notation introduced in the lecture and also choose our domain accordingly with  $\Omega = [0.0, 1.0]^2$ .

- In the following image you see an incomplete regular sparse grid  $V_2^1$ . Insert the missing grid points using small **squares**. What are the level-index-vector pairs  $\vec{l}, \vec{i}$  for each of them?



- Use the (modified) picture from the previous task to perform two steps of adaptive refinement:
  - Refine grid point  $\vec{l}, \vec{i} = (1, 2), (1, 3)$ : create all hierarchical children. Draw its children as small **triangles**. Make sure that you also insert all missing hierarchical parents (and parents of parents, ...) of these children to make the grid suitable for typical algorithms on sparse grids.

- (b) Now refine grid point  $(2, 2), (3, 3)$ . Again, do not forget to create all missing parents. Draw all new points as small **crosses**.

## 2 Orthogonality

We first consider orthogonality of functions  $[a, b] \rightarrow \mathbb{R}$  in the two scalar products we already know

- $L^2$  scalar product:

$$(u, v)_2 := \int_a^b u(x)v(x) dx$$

- “energy scalar product”:

$$(u, v)_a := \int_a^b u'(x)v'(x) dx,$$

We assume that the space of functions under consideration again be well-defined such that  $(u, u) > 0$  for  $u \neq 0$  is ensured.

- (i) Show that for  $g_k : [0, 2\pi] \rightarrow \mathbb{R}, g_k(x) = \sin(kx)$  and  $k, j \in \mathbb{N}$  the  $L_2$  scalar product is

$$(g_k, g_j)_2 = \begin{cases} 0 & \text{for } k \neq j, \\ \pi & \text{else.} \end{cases}$$

- (ii) Which functions of the hierarchical basis are orthogonal to each other w.r.t. the  $L_2$  scalar product? What about the energy scalar product?
- (iii) Let  $V$  be a vector space with  $\dim V = n < \infty$  with scalar product  $(\cdot, \cdot)$  and associated norm  $\|x\| := \sqrt{(x, x)}$ . Also let  $\Psi = \{\psi_1, \dots, \psi_n\} \subset V$  a orthonormal system, i.e.

$$(\psi_i, \psi_j) = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{else.} \end{cases}$$

- a) Show that for

$$x = \sum_{i=1}^n \alpha_i \psi_i$$

the following holds:

$$\|x\| = \sqrt{\sum_{i=1}^n \alpha_i^2}.$$

- b) Show that  $\Psi$  is a linearly independent system!
- c) Show for every  $x \in V$ :

$$x = \sum_{i=1}^n (x, \psi_i) \psi_i.$$