

Mock Exam Algorithms of Scientific Computing (Univ.-Prof. Dr. Michael Bader) SS 2017	Page 1/13
Last name, first name, student ID:	Signature:

## General Instructions

### Material:

You may only use one hand-written sheet of paper (size A4, on both pages).

Any other material including electronic devices of any kind is forbidden.

Use only the exam paper that was handed out to solve the exercises. If the space on a page is not enough, choose any empty space provided and state where you continue with your solution. For additional notes and sketches, you can obtain separate exam sheets.

Do not use pencil, or red or green ink.

### Errors and ambiguities:

If you think that a question contains an error or ambiguity, then correct it or choose an interpretation that allows you to complete the exercise.

Make sure to point out your correction or disambiguation in your solution!

Note that questions, esp. concerning errors or ambiguities, will not be answered during the exam.

### General hint:

Often, exercises b), c), etc. can be solved without the results from the previous exercise a); if you are stuck with exercise a), then don't immediately skip exercises b), c), etc.

### Maximum score:

The total number of points for all exercises exceeds 40, however grades will be computed relative to a maximum score of 40 points.

### Working time:

100 minutes.

**Please switch off your cell phones!**

## Proposed Solution

1	2	3	4	$\Sigma$
/11	/12	/10	/11	/44

Last name, first name, student ID:

## 1 Quarter-Wave Discrete Sine Transform ( $\approx 5 + 3 + 3 = 11$ points)

Given is a real-valued input data-set  $f_0, \dots, f_{2N-1} \in \mathbb{R}$  (i.e., length  $2N$ ) with the symmetry conditions  $f_{2N-n-1} = -f_n$ . Complete the tasks below.

a) Show that the corresponding Quarter-Wave Fourier coefficients

$$F_k = \frac{1}{2N} \sum_{n=0}^{2N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} \quad (1)$$

have only imaginary values and can be written as

$$F_k = -\frac{i}{N} \sum_{n=0}^{N-1} f_n \sin\left(\frac{\pi k}{N} \left(n + \frac{1}{2}\right)\right) \quad (2)$$

The proof is done in the following steps:

- ① Isolate the symmetry condition
- ② Insert the symmetry condition
- ③ Assemble terms to a sum over  $f_n$
- ④ Make terms “imaginary”

Last name, first name, student ID:

$$\begin{aligned}
F_k &= \frac{1}{2N} \sum_{n=0}^{2N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} \\
&\stackrel{\textcircled{1}}{=} \frac{1}{2N} \left( \sum_{n=0}^{N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} + \sum_{n=N}^{2N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} \right) \\
&\stackrel{\textcircled{2}}{=} \frac{1}{2N} \left( \sum_{n=0}^{N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} - \sum_{n=N}^{2N-1} f_{2N-n-1} \omega_{2N}^{-k(n+\frac{1}{2})} \right) \\
&= \frac{1}{2N} \left( \sum_{n=0}^{N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} - \sum_{n=0}^{N-1} f_n \omega_{2N}^{-k(2N-n-1+\frac{1}{2})} \right) \\
&= \frac{1}{2N} \left( \sum_{n=0}^{N-1} f_n \omega_{2N}^{-k(n+\frac{1}{2})} - \sum_{n=0}^{N-1} f_n \omega_{2N}^{k(n+\frac{1}{2})} \right) \\
&\stackrel{\textcircled{3}}{=} \frac{1}{2N} \sum_{n=0}^{N-1} f_n \underbrace{\left( \omega_{2N}^{-k(n+\frac{1}{2})} - \omega_{2N}^{k(n+\frac{1}{2})} \right)}_{= \omega_{2N}^{-k(n+\frac{1}{2})} - \left( \omega_{2N}^{k(n+\frac{1}{2})} \right)^* = -2i \operatorname{Im}\{\omega_{2N}^{-k(n+\frac{1}{2})}\}} \\
&\stackrel{\textcircled{4}}{=} \frac{-2i}{2N} \sum_{n=0}^{N-1} f_n \underbrace{\operatorname{Im}\{e^{i2\pi k(n+\frac{1}{2})/2N}\}}_{= \operatorname{Im}\left\{ \cos\left(\frac{\pi k(n+\frac{1}{2})}{N}\right) + i \sin\left(\frac{\pi k(n+\frac{1}{2})}{N}\right) \right\}} \\
&= -\frac{i}{N} \sum_{n=0}^{N-1} f_n \sin\left(\frac{\pi k}{N} \left(n + \frac{1}{2}\right)\right) \in \mathbb{C} \quad \text{q.e.d.}
\end{aligned}$$

Last name, first name, student ID:

b) Show that the coefficients  $F_k$  of the QW-DST again justify a symmetry condition!

Since there are only sine terms, we assume (and get) an odd symmetry:

$$\begin{aligned}
 F_{2N-k} &= -\frac{i}{N} \sum_{n=0}^{N-1} f_n \sin \left( \frac{\pi}{N} (2N - k) \left( n + \frac{1}{2} \right) \right) \\
 &= -\frac{i}{N} \sum_{n=0}^{N-1} f_n \sin \left( 2\pi + \frac{\pi k}{N} \left( n + \frac{1}{2} \right) \right) \\
 &= \frac{i}{N} \sum_{n=0}^{N-1} f_n \sin \left( \frac{\pi k}{N} \left( n + \frac{1}{2} \right) \right) \\
 &= F_k
 \end{aligned}$$

Since all  $F_k = F_{2N-k}$ , we need the  $F_k$  only for  $k = 0, \dots, N$  for a Sine Transform.

Last name, first name, student ID:

- c) Let  $\text{real-FFT}(g, N)$  be a procedure that computes the Fourier coefficients  $G_k$  efficiently from a real data-set  $g$  that consists of  $2N$  values  $g_n$ .

Describe a **procedure**  $\text{QW-DST}(g, N)$  that uses the given procedure  $\text{real-FFT}$  to compute the coefficients  $F_k$  for  $k = 0, \dots, N-1$  from equation (2) for the (non-symmetrical) real data  $f_0, \dots, f_{N-1}$ , stored in the parameter field  $g$ .

Note:  $\text{real-FFT}(g, N)$  **does not** compute a QW-RDFT!

### QW-DST algorithm reduced to real FFT:

- (a) For  $n = 0, \dots, N-1$  set

$$g_n = f_n,$$

$$g_{2N-n-1} = -f_n.$$

- (b) Use  $2N$   $\text{real-FFT}$  to compute  $G_k$  from  $g_n$  (for  $k = 0, \dots, N$ ).

- (c) Calculate the QW-DST coefficients  $F_k = G_k \omega_{2N}^{-\frac{k}{2}}$

Last name, first name, student ID:

## 2 Wavelets Approximation

( $\approx 8 + 4 = 12$  points)

The cubic Battle-Lemarié wavelet is given by the coefficients  $\{c_k\}$ :

$$c_0 = \frac{1}{8} \quad c_1 = \frac{1}{2} \quad c_2 = \frac{3}{4} \quad c_3 = \frac{1}{2} \quad c_4 = \frac{1}{8}. \quad (3)$$

Starting with the mother hat function

$$\phi_0(t) = \max\{1 - |t|, 0\}, \quad (4)$$

we can compute an approximation of the scaling and wavelet functions by iterating over their dilation equations.

Ignoring the scaling factor such as  $\frac{1}{\sqrt{2}}$ , the dilation equation for the scaling (father) function is given by

$$\phi_{n+1}(t) = \sum_{k=0}^4 c_k \cdot \phi_n(2t - k), \quad (5)$$


and the dilation equation for the wavelet (mother) function is given by

$$\psi_{n+1}(t) = \sum_{k=0}^4 (-1)^k c_{K-k} \cdot \phi_n(2t - k), \quad \text{where } K = 4. \quad (6)$$

- a) In the given figure below, sketch the first approximation of the scaling function  $\phi_1(t)$  and the wavelet function  $\psi_1(t)$  on the interval  $[-1, 3]$ .

**Hint 1:** Use the given helper table on the right to compute the values of  $\phi_1(t)$  and  $\psi_1(t)$  at given points to help for your sketching.


**Hint 2:** Backup figure and helper table is provided in case you need to correct your solutions.



	$\phi_1(t)$	$\psi_1(t)$
$t = -1.0$	0.0	0.0
$t = -0.5$	0.0	0.0
$t = 0.0$	0.125	0.125
$t = 0.5$	0.5	-0.5
$t = 1.0$	0.75	0.75
$t = 1.5$	0.5	-0.5
$t = 2.0$	0.125	0.125
$t = 2.5$	0.0	0.0
$t = 3.0$	0.0	0.0

Last name, first name, student ID:

**Backup figure and helper table**



	$\phi_1(t)$	$\psi_1(t)$
$t = -1.0$		
$t = -0.5$		
$t = 0.0$		
$t = 0.5$		
$t = 1.0$		
$t = 1.5$		
$t = 2.0$		
$t = 2.5$		
$t = 3.0$		

Last name, first name, student ID:

b) By observation, what conclusion can you draw for the integral of the obtained wavelet, i.e.,  $\int_{-\infty}^{+\infty} \psi_1(t)$ ?

Explain, what this implies for wavelets in general.

### Observation:

- Area above x-axis equals area below x-axis,  
i.e.,  $\rightarrow \int_{-\infty}^{+\infty} \psi_1(t) = 0$

### What it implies:

- The mean is zero  $\rightarrow$  The wavelet basis functions will represent fluctuations to the average of the coarser level.
- (admissibility condition is met)



### 3 Sparse Grids

( $\approx 6 + 4 = 10$  points)

a) Name **two** different data structures for sparse grids. For each of them, discuss the following points:

- (i) Hierarchization/dehierarchization (Hint: consider data access and traversal complexity)
- (ii) Spatial adaptivity
- (iii) Memory consumption

Discuss any two of these data structures:

#### Arrays

- (i) Access by index  $O(1)$ , need mapping between hierarchical index  $(\vec{l}, \vec{i})$  and “flat“ (1D) index  $j$ . Hierarchical neighbors can be deduced from the current node’s hierarchical index  $(\vec{l}, \vec{i})$ .
- (ii) Cannot add or delete element  $\rightarrow$  Does not support spatial adaptivity. However, dimensional adaptivity can be accommodated.
- (iii) Store only data, i.e.,  $v_j$ , at each grid point. No need to store  $(\vec{l}, \vec{i})$  if correct mapping is provided. Cache efficient!

#### Trees


- (i) One hierarchical neighbour is the parent node, the other one (ancestor) can be found from the root node. The neighbours can be passed as additional parameters in the algorithm.
- (ii) Spatial Adaptivity is done straightforward by adding the nodes corresponding to the adaptive points.
- (iii) On each node:  $2d$  pointers to children,  $d$  optional pointers for parent (ancestors can be trace back through parent pointer)  $\rightarrow$  relatively high memory consumption.

#### Hash tables


- (i) Access by Key—hierarchical index  $(\vec{l}, \vec{i}) \rightarrow$  complexity  $O(1)$ , computation of indices (hashing).
- (ii) Spatial Adaptivity poses no problem, additional points can be inserted independently of existing points in the data structure.
- (iii) Memory consumption is low, because for each node, we only need to store a Key  $(\vec{l}, \vec{i})$  and a data  $v_j$ . Cache inefficient due to hashing.

Last name, first name, student ID:

- b) Discuss whether the following two grids are valid sparse grids. Explain your reasons. You can annotate the grids directly.



Valid: adaptive refinement, no missing parent



Invalid: point in red missing a parent (blue)



Last name, first name, student ID:

$$\bullet D \longrightarrow C \downarrow C \downarrow D \leftarrow D \leftarrow D \uparrow B \rightarrow A \uparrow A \leftarrow D$$

b) Arithmetical representation of the Peano-Meander curve

1) Given is a parametrization  $q(t)$  of the Peano-Meander curve

$$q(0_9.n_1n_2n_3n_4\dots) = Q_{n_1} \circ Q_{n_2} \circ Q_{n_3} \circ Q_{n_4} \circ \dots \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (7)$$

where  $t = 0_9.n_1n_2n_3n_4\dots$  is the representation of  $t$  in a base nine system. Determine the operators  $Q_1$ ,  $Q_4$  and  $Q_6$ .

$$\begin{aligned} Q_1 \begin{pmatrix} x \\ y \end{pmatrix} &= \frac{1}{3} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \frac{1}{3} \\ 0 \end{pmatrix} \\ &= \frac{1}{3} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \frac{1}{3} \\ 0 \end{pmatrix} \end{aligned}$$

$$Q_4 \begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \frac{2}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{pmatrix}$$

$$\begin{aligned} Q_6 \begin{pmatrix} x \\ y \end{pmatrix} &= \frac{1}{3} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \frac{2}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{pmatrix} \\ &= \frac{1}{3} \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} \frac{2}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{pmatrix} \end{aligned}$$

2) Compute the coordinates of  $q(\frac{2}{3})$  and  $q(\frac{1}{2})$ .

$$\frac{2}{3} = 0_9.6$$

$$q\left(\frac{2}{3}\right) = Q_6 \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{2}{3} \\ \frac{2}{3} \\ \frac{2}{3} \end{pmatrix}$$

$$\frac{1}{2} = 0_9.44444\dots$$

Last name, first name, student ID:

$$q\left(\frac{1}{2}\right) = Q_4 \circ Q_4 \circ \dots \left(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}\right) = \lim_{n \rightarrow \infty} Q_4^n \left(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}\right)$$

Write  $Q_4$  in the following form:

$$Q_4 = \underbrace{\begin{pmatrix} \frac{1}{3} & 0 \\ 0 & \frac{1}{3} \end{pmatrix}}_{=:A} \underbrace{\begin{pmatrix} x \\ y \end{pmatrix}}_{=:v} + \underbrace{\begin{pmatrix} \frac{2}{3} \\ \frac{2}{3} \end{pmatrix}}_{=:b} = Av + b.$$

From this, we get

$$Q_4^n v = A^n v + A^{n-1} b + \dots + Ab + b$$

and

$$(I - A)(A^{n-1} b + \dots + Ab + b) = b - A^n b = (I - A^n) b$$

Now we can compute

$$\begin{pmatrix} \alpha \\ \beta \end{pmatrix} =: \lim_{n \rightarrow \infty} Q_4^n \left(\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}\right) = \lim_{n \rightarrow \infty} \left( \underbrace{A^n v}_{\rightarrow 0} (I - A)^{-1} (I - \underbrace{A^n}_{\rightarrow 0}) b \right) = (I - A)^{-1} b$$

$$\Leftrightarrow (I - A) \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = b \Leftrightarrow \begin{pmatrix} \frac{2}{3} & 0 \\ 0 & \frac{2}{3} \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \begin{pmatrix} \frac{2}{3} \\ \frac{2}{3} \end{pmatrix}$$

**Solution:**  $\alpha = 1$ ,  $\beta = 1$ .